

# WHERE ARE YOU FOCUSED?



Multifocal IOLs

**LIVE THE LIFE  
YOU WANT TO SEE.**

**TECNIS<sup>®</sup>**  
Multifocal IOL

**CAUTION:** Federal law restricts this device to sale, distribution and use by or on the order of a physician.

*See reverse for continued Indications and Important Safety Information.*

*Johnson & Johnson* VISION

# EMPOWER WHAT MAKES LIFE YOURS.



Presbyopia is a gradual, natural blurring of sight at close distances that occurs as the lens of your eye loses flexibility. You might already know it as the reason you wear reading glasses. **But here's something you may not know: cataract surgery can be a great opportunity to reduce your dependence on glasses.**



If you have both presbyopia (with or without astigmatism) and cataracts, you may have the opportunity to address them simultaneously with a multifocal IOL.

# SEE LIFE YOUR WAY.

Cataract surgery is trading one thing for another. When your cataract is removed, it is replaced by a small, soft lens called an *intraocular lens* — or *IOL* for short. You won't be able to feel or see the IOL, but it will stay inside your eye to help you see clearly.

**Multifocal IOLs address presbyopia as part of cataract surgery. If you have astigmatism, there are Multifocal Toric IOLs to address your vision needs.**

**TECNIS®**  
Multifocal IOL

**TECNIS®**  
Multifocal Toric II IOL

**TECNIS®**  
Multifocal Toric II IOL

+2.75 D  
Toric II

+3.25 D  
Toric II

**TECNIS®** Multifocal IOLs give you high-quality distance vision and personalized near vision for clear focus where you need it most.



## PRESBYOPIA

Age

**40**

Typically becomes noticeable around the age of 40<sup>1</sup>



Makes it difficult to focus on things up close (near vision)



Can be addressed with a multifocal or extended depth of focus IOL



Other solutions include reading glasses, contacts or an additional surgery



### With Presbyopia

*These images are for illustrative purposes only and do not represent actual data derived from studies. These illustrative simulations are intended to help you better understand your vision in certain eye conditions. [See more at tecnisvisionsimulator.com](http://See more at tecnisvisionsimulator.com).*



Help you see far away



Personalize your vision  
between 13 and 20 inches  
to your unique needs



Mean you may wear  
glasses less often overall



Ask your doctor about other IOL  
opportunities — including astigmatism  
management and continuous vision  
across the full visual range.



### Without Presbyopia

**Talk to your doctor about the  
ways you read, work and play,  
and find out if TECNIS® Multifocal  
IOLs are right for you.**

**TECNIS**<sup>®</sup>  
Multifocal IOL

**TECNIS**<sup>®</sup>  
Multifocal Toric II IOL

**TECNIS**<sup>®</sup>  
Multifocal Toric II IOL

+2.75 D

Toric II

+3.25 D

Toric II

## INDICATIONS AND IMPORTANT SAFETY INFORMATION FOR THE TECNIS<sup>®</sup> MULTIFOCAL 1-PIECE AND TECNIS<sup>®</sup> MULTIFOCAL TORIC II IOLS (Continued)

**INDICATIONS:** The TECNIS<sup>®</sup> Multifocal 1-Piece intraocular lenses are indicated for primary implantation for the visual correction of aphakia in adult patients with and without presbyopia in whom a cataractous lens has been removed by phacoemulsification and who desire near, intermediate, and distance vision with increased spectacle independence. The TECNIS<sup>®</sup> Multifocal Toric II lens models ZKU150, ZKU225, ZKU300, ZKU375 and ZLU150, ZLU225, ZLU300, ZLU375 are indicated for primary implantation for the visual correction of aphakia and for reduction of residual refractive astigmatism in adult patients with or without presbyopia, with greater than or equal to 1 diopter of preexisting corneal astigmatism, in whom a cataractous lens has been removed in order to provide near, intermediate and distance vision. The IOLs are intended for capsular bag placement only.

**RISKS:** Whereas your use of glasses may decrease, it may come at the cost of some sharpness of your vision. You may also notice more halos (rings of light around a light source) or glare (scattered light effect around a light source). Complications with cataract surgery are rare and may include the worsening of your vision, bleeding, or infection. Call your eye doctor right away if you experience any itching, pain, flashing lights, “floaters”, redness, severe headache, upset stomach, light sensitivity, or watery eyes after surgery.

**WARNINGS:** Less than 1% of patients may not be satisfied and request to remove the IOL. Children under the age of 2 those with previous trauma to their eye should not receive this device. Existing eye conditions and surgical difficulties may increase the risk of complications. Your doctor will determine if you are a good candidate for lens replacement. You may experience vision problems or distortions and may require a second surgery to have your doctor correctly reposition the lens.

**PRECAUTIONS:** If your eye is not healthy (including glaucoma) you may not get the full benefits of the TECNIS<sup>®</sup> Multifocal IOL. Your vision with a multifocal IOL may not be good enough to perform very near or detailed “up-close” work without glasses. There is a chance that you still may need glasses for distance vision with a Toric IOL.

**SIDE EFFECTS:** Multifocal IOL implants may be inadvisable in patients with conditions such as macular degeneration, retinal pigment epithelium changes, and glaucoma. In rare cases, severe incidents of halos and/or glare may result in the multifocal IOL being explanted and replaced with another type of IOL. Side effects of cataract surgery with the TECNIS<sup>®</sup> Multifocal Toric II IOLs include the need for a second surgery to reposition the lens or repair the retina, swelling/thickening of the retina and separation of the retina from the surrounding tissues.

**ATTENTION:** Please see Directions for Use for a complete list of Indications and Important Safety Information

### Reference

1. Presbyopia. American Optometric Association Web site. <https://www.aoa.org/patients-and-public/eye-and-vision-problems/glossary-of-eye-and-vision-conditions/presbyopia>. Accessed October 28, 2017.